



PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM	Books
Search	PubMed	<input type="text"/>	for	<input type="text"/>	Go	Clear		
<input checked="" type="checkbox"/> Limits		Preview/Index		History	Clipboard	Details		

Display	Abstract	<input type="text"/>	Show:	20	<input type="text"/>	Sort	<input type="text"/>	Send to	Text	<input checked="" type="checkbox"/>
---------	----------	----------------------	-------	----	----------------------	------	----------------------	---------	------	-------------------------------------

☐ 1: FEBS Lett. 1997 Jun 9;409(2):312-6.

[Related Articles, Links](#)

Entrez
PubMed

ELSEVIER SCIENCE
FULL-TEXT ARTICLE

The nodulation gene nolK of Azorhizobium caulinodans is involved in the formation of GDP-fucose from GDP-mannose.

Mergaert P, Van Montagu M, Holsters M.

PubMed
Services

Department of Genetics, Flanders Interuniversity Institute for Biotechnology (VIB), Universiteit Gent, Belgium.

Related
Resources

The nolK gene of *Azorhizobium caulinodans* is essential for the incorporation of a fucosyl group in Nod factors. A NAD(P)-binding site is present in the NolK amino acid sequence and the gene is homologous to *Escherichia coli* genes, presumably involved in GDP-fucose synthesis. Protein extracts of *A. caulinodans*, overexpressing nolK, have an enzyme activity that synthesizes GDP-fucose from GDP-mannose. nolK most probably encodes a 4-reductase performing the last step in GDP-fucose synthesis. Wild-type *A. caulinodans* produces a population of fucosylated and non-fucosylated molecules but the nolK-overexpressing strain produces only fucosylated Nod factors. Thus, the production of activated fucosyl donors is a rate-limiting step in Nod factor fucosylation.

PMID: 9202168 [PubMed - indexed for MEDLINE]

Display	Abstract	<input type="text"/>	Show:	20	<input type="text"/>	Sort	<input type="text"/>	Send to	Text	<input checked="" type="checkbox"/>
---------	----------	----------------------	-------	----	----------------------	------	----------------------	---------	------	-------------------------------------

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

[Department of Health & Human Services](#)

[Freedom of Information Act](#) | [Disclaimer](#)

WEST

Help

Logout

Interrupt

Main Menu

Search Form

Posting Counts

Show S Numbers

Edit S Numbers

Preferences

Cases

Search Results -

Terms	Documents
L4 same GDP-4-keto-6-deoxy-D-mannose	1

Database:

US Patents Full-Text Database
 US Pre-Grant Publication Full-Text Database
 JPO Abstracts Database
 EPO Abstracts Database
 Derwent World Patents Index
 IBM Technical Disclosure Bulletins

Search:

L6

Refine Search

Recall Text

Clear

Search HistoryDATE: Saturday, June 07, 2003 [Printable Copy](#) [Create Case](#)**Set Name Query**

side by side

Hit Count Set Name

result set

DB=USPT,PGPB,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ

<u>L6</u>	L4 same GDP-4-keto-6-deoxy-D-mannose	1	<u>L6</u>
<u>L5</u>	L4 same D-mannose	1	<u>L5</u>
<u>L4</u>	L3 same (arabidopsis or thaliana)	236	<u>L4</u>
<u>L3</u>	epimerase or reductase	23381	<u>L3</u>
<u>L2</u>	GDP-4-keto-6-deoxy-D-mannose 3,5-epimerase-4-reductase	0	<u>L2</u>
<u>L1</u>	epimerase-4-reductase	1	<u>L1</u>

END OF SEARCH HISTORY

WEST**End of Result Set**

Generate Collection

Print

L6: Entry 1 of 1

File: PGPB

Jan 31, 2002

DOCUMENT-IDENTIFIER: US 20020012979 A1

TITLE: VITAMIN C PRODUCTION IN MICROORGANISMS AND PLANTS

Detail Description Paragraph (43):

[0074] The ability to substitute GDP-4-keto-6-D-mannose epimerase/reductase for GDP-D-mannose:GDP-L-galactose epimerase to enhance L-ascorbic acid biosynthesis in plants or microorganisms depends on the ability of GDP-4-keto-6-deoxy-D-mannose epimerase/reductase to act directly on GDP-D-mannose to form GDP-L-galactose. Evidence supporting this possibility already exists. Arabidopsis thaliana murl mutants are defective in GDP-D-mannose-4,6-dehydratase activity (Bonin, et al., 1997, Proc. Natl. Acad. Sci. 94:2085-2090). These mutants are thus blocked in GDP-L-fucose biosynthesis, and consequently have less than 2% of the normal amounts of L-fucose in the primary cell walls of aerial portions of the plant (Zablackis, et al., 1996, Science 272:1808-1810). The murl mutants are more brittle than wild-type plants, are slightly dwarfed and have an apparently normal life cycle (Zablackis, et al., 272:1808-1810). When murl mutants are grown in the presence of exogenous L-fucose, the L-fucose content in the plant is restored to the wild-type state (Bonin, et al., 1997, Proc. Natl. Acad. Sci. 94:2085-2090). It was discovered (Zablackis, et al., 1996, Science 272:1808-1810) that murl mutants contain, in the hemicellulose xyloglucan component of the primary cell wall, L-galactose in place of the normal L-fucose. L-galactose is not normally found in the xyloglucan component, but in murl mutants L-galactose partly replaces the terminal L-fucosyl residue. Bonin, et al. (1997, Proc. Natl. Acad. Sci. 94:2085-2090) hypothesized that in the absence of a functional GDP-D-mannose-4,6-dehydratase in the murl mutants, the GDP-4-keto-6-deoxy-D-mannose epimerase/reductase normally involved in L-fucose synthesis may be able to use GDP-D-mannose directly, forming GDP-L-galactose. Another possibility, however, is that the enzymes involved in L-ascorbic acid biosynthesis in A. thaliana are responsible for forming GDP-L-galactose in the murl mutant. If this were true, it would suggest that in the wild-type plant, some mechanism exists that prevents GDP-L-galactose formed in the L-ascorbic acid pathway from entering cell wall biosynthesis and substituting for (competing with) GDP-L-fucose for incorporation into the xyloglucan component (since L-galactose is not present in the primary cell wall of the wild-type plant).

Detail Description Paragraph (69):

[0100] According to the present invention, a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase can be a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase from any organism, including Arabidopsis thaliana, Escherichia coli, and human. A nucleic acid sequence encoding a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase from Arabidopsis thaliana is represented herein by SEQ ID NO:1. SEQ ID NO:1 encodes a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase having an amino acid sequence represented herein as SEQ ID NO:2. A nucleic acid sequence encoding a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase from Escherichia coli is represented herein by SEQ ID NO:3. SEQ ID NO:3 encodes a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase having an amino acid sequence represented herein as SEQ ID NO:4. A nucleic acid sequence encoding a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase from homo sapiens is represented herein by SEQ ID NO:5. SEQ ID NO:5 encodes a GDP-4-keto-6-deoxy-D-mannose epimerase/reductase having an amino acid sequence represented herein as SEQ ID NO:6.